

MIT/LCS/TM-189

PERSISTENCE OF VECTOR REPLACEMENT  
SYSTEMS IS DECIDABLE

Ernst Mayr

January 1981

# Persistence of Vector Replacement Systems is Decidable

*by*

Ernst Mayr  
Visiting Scientist  
Laboratory for Computer Science  
Massachusetts Institute of Technology

December 1, 1980

## **Abstract:**

In a persistent vector replacement system (VRS) or Petri net, an enabled transition can become disabled only by firing itself. Here, an algorithm is presented which allows to decide whether an arbitrary VRS is persistent or not, and if so, to construct a semilinear representation of the set of states reachable in the system.

**Key Words and Phrases:** Vector Replacement System, Petri Net, Persistence, Representation of Reachability Set

This work was supported in part by the Deutsche Forschungsgemeinschaft, Grant No. I3 Ma 870/1-1, and in part by the National Science Foundation, Grant No. MCS 8010707

Journal of Environmental Systems Management  
Volume 1, Number 1  
1980

1980, Volume 1, Number 1

Abstracts

The following abstracts are from the first issue of the Journal. The abstracts are arranged in alphabetical order of the author's name. The abstracts are intended to provide a brief summary of the articles in this issue. The abstracts are not intended to be a substitute for the full text of the articles.

1. A. J. ... (Title of article 1)

2. B. C. ... (Title of article 2)

## 1. Introduction

In this report, an effective algorithm is presented which for any given arbitrary *vector replacement system* (VRS) [8] allows to decide whether it is persistent or not. This algorithm is an extension of the one presented in [11] which relies on the persistence of the given Petri net when it constructs a finite semilinear representation of its reachability set. The algorithm here is self-contained, based on a recursive construction of semilinear representations for subsets of the counter set of the VRS. If the VRS is persistent, the whole counter set is obtained by the algorithm, thus also giving a semilinear representation of the reachability set. For further motivation to study persistent systems the reader is referred to [6,10,13,17]. In [5], it is proved that the decision problem for the persistence of one transition in a VRS or Petri net is recursively equivalent to the decidability of the reachability problem, but it is also conjectured that the persistence of a VRS or Petri net can be decided completely independent from the reachability problem. The algorithm presented in the sequel does not rely on an algorithm for the general reachability problem. Throughout the paper, VRS terminology is used; the transition to and from Petri nets is straightforward [4].

## 2. Notation and preliminaries

A *Vector Replacement System* (VRS) [8] is a pair  $(T, m)$  where  $T = \{t_1, \dots, t_w\}$  is a finite set of *transitions*  $t_i = (u_i, v_i) \in \mathbb{N}^v \times \mathbb{Z}^w$  with  $u_i + v_i \geq 0$  and  $m \in \bar{\mathbb{N}}^v$  is the *initial vector* ( $v \in \mathbb{N}$ ).  $t_i \in T$  is *applicable* at  $m' \in \bar{\mathbb{N}}^v$  iff  $u_i \leq m'$  (written  $a(t_i, m')$ ), application of  $t_i$  at  $m'$  takes  $m'$  to  $m' + v_i$  (written  $m' \xrightarrow{t_i} m' + v_i$ ).

For  $\tau = t_{i_1} \dots t_{i_r} \in T^*$  we define inductively

- i)  $a(\tau, m') := r = 0 \vee a(t_{i_1}, m') \wedge a(t_{i_2} \dots t_{i_r}, m' + v_{i_1})$ ;
- ii)  $\delta\tau := \sum_{j=1}^r v_{i_j}$ ;
- iii)  $m' \xrightarrow{\tau} m'' := a(\tau, m') \wedge (m'' = m' + \delta\tau)$ .

The *reachability set*  $R(T, m)$  of  $(T, m)$  is  $R(T, m) := \{m'; (\exists \tau \in T^*) [m \xrightarrow{\tau} m']\}$ . Let  $\Phi: T^* \rightarrow \mathbb{N}^w$  denote the Parikh mapping. The *counter set*  $C(T, m)$  of  $(T, m)$  is given by

$$C(T, m) := \{\Phi(\tau); \tau \in T^* \wedge a(\tau, m)\}.$$

Let  $V \in \mathbb{Z}^{v,w}$  be the integer matrix whose  $i$ -th column is  $v_i$ ,  $i \in I_w$ . Obviously, we have

- i)  $(\forall \tau \in T^*) [\delta\tau = V\Phi(\tau)]$ ;
- ii)  $R(T, m) = \{m + Vc; c \in C(T, m)\}$ .

A *linear set*  $L \subseteq \mathbb{N}^w$  is a set of the form  $L = \{a + \sum_{i=1}^r n_i b_i; (n_1, \dots, n_r) \in \mathbb{N}^r\}$  for some  $r \in \mathbb{N}$ ,  $a, b_1, \dots, b_r \in \mathbb{N}^w$ . A *semilinear set* is a finite union of linear sets. Semilinear sets are exactly those sets definable by expressions in Presburger Arithmetic, i.e. the first order theory of the nonnegative integers with addition [16]. There is an effective procedure to construct semilinear representations of the sets defined by Presburger expressions [3,14].

### Definition 1:

A vector  $m' \in \bar{\mathbb{N}}^v$  is *persistent* iff

$$(\forall t_i, t_j \in T) [(i \neq j) \wedge a(t_i, m') \wedge a(t_j, m')] \Rightarrow a(t_i t_j, m').$$

$(T, m)$  is *persistent* iff all  $m' \in R(T, m)$  are persistent.

It is known that for a persistent VRS  $(T, m)$ ,  $R(T, m)$  and  $C(T, m)$  are effectively constructible semilinear sets [9,12].

The following algorithm for the construction of the *reachability graph*  $RG(T, m)$  works for arbitrary VRS's, it does not assume persistence. In this algorithm, which is a slight modification of one originally given in [6], a digraph with labelled nodes and edges is

\* $\mathbb{N}$  denotes the set of nonnegative integers,  $\mathbb{Z}$  the set of integers, and  $\bar{\mathbb{N}} := \mathbb{N} \cup \{\omega\}$  the set  $\mathbb{N}$  augmented by the "infinite" number  $\omega$  with  $\pm n + \omega = \omega \pm n = \omega$  and  $n < \omega$  for all  $n \in \mathbb{N}$ . For  $i \in \mathbb{N}$ ,  $I_i$  stands for the set  $\{1, \dots, i\}$ .

constructed: The label  $t(e)$  of an edge  $e$  is an element of  $T$ , and each node  $k$  obtains a label  $\bar{m}(k) \in \bar{N}^t$ . (In pictures, parallel edges are merged into one which receives the union of the edge labels.) From these labels, marks have to be distinguished which in the algorithm serve to decide which nodes still have to be dealt with.

**Algorithm 1:**

```

begin
  start with an unmarked node  $r$  (the "root") with  $\bar{m}(r) := m$ ;
  while there is an unmarked node do
    select nondeterministically an unmarked node  $k$ ;
    mark  $k$ ;
    for all  $j \in I_\omega$  with  $a(t_j, \bar{m}(k))$  do
      add to the graph constructed so far a new unmarked node  $k'$  and an edge  $e$ 
      from  $k$  to  $k'$  with  $t(e) := t_j$ ;
      for  $i := 1, \dots, v$  do
        if there is a node  $k''$  on a (not necessarily simple) path from  $r$  to  $k$  with
            $\bar{m}(k'') \leq \bar{m}(k) + v_j$  and  $(\bar{m}(k''))_i < (\bar{m}(k) + v_j)_i$ 
        then
           $(\bar{m}(k'))_i := \omega$ 
        else
           $(\bar{m}(k'))_i := (\bar{m}(k) + v_j)_i$ 
        fi
      od;
      if there is a node  $k'' \neq k'$  in the graph constructed so far with  $\bar{m}(k'') = \bar{m}(k')$ 
      then
        identify  $k'$  with this  $k''$ 
      fi
    od
  od
end Algorithm 1.

```

Note that new  $\omega$ -components are introduced independently for different coordinates as for each coordinate  $i$  a different node  $k''$  may be found.


The proof of the termination of Algorithm 1 is very similar to the one given in [6] and won't be presented here.

**Example:**

For the persistent VRS  $(T, m)$  with

$$T = \{(0^6, 10^5), (10^5, 012100 - 10^5), (0210^3, 0^410 - 0210^3), (0^21^20^2, 0^5 - 0^21^20^2), (0^41^2, 10^5 - 0^41^2)\}$$

and  $m = 10^5$  (which is short for  $(1, 0, 0, 0, 0, 0) \in \bar{\mathbb{N}}^6$ ) some run of Algorithm 1 produces the graph  $RG(T, m)$ :



**Lemma 1:**

Given  $(T, m)$ , the set  $NPC(T, m) := \{c \in \mathbb{N}^w; m + Vc \text{ is } \geq 0 \text{ and not persistent}\}$  is an effectively constructable semilinear set.

*Proof:*

By Definition 1,  $NPC(T, m)$  equals the set

$$\{c \in \mathbb{N}^w; (\exists t_i, t_j \in T)[i \neq j \wedge m + Vc \geq u_i \wedge m + Vc \geq u_j \wedge m + Vc + v_i \not\geq u_j]\}.$$

Hence,  $NPC(T, m)$  can be defined in Presburger Arithmetic, and a semilinear representation can effectively be found. ■

### 3. Some properties of $RG(T, m)$

Let  $k$  be a node in  $RG(T, m)$  for some arbitrary VRS  $(T, m)$ .

#### Lemma 2:

The sets

$CT_k := \{\Phi(\tau); \tau \text{ is the edge-labelling sequence of a path in } RG(T, m) \text{ from } k \text{ to } k\}$ , and

$CT_k^+ := \{c \in CT_k; \forall c \geq 0\}$

are effectively constructable semilinear sets.

#### Proof:

Regarding the strongly connected component (SCC) of  $k$  in  $RG(T, m)$ , stripped of the node marking  $\bar{m}$ , as the transition diagram of a finite automaton over  $T$ , the set  $CT_k$  corresponds to the Parikh image of the regular language accepted by that finite automaton with  $k$  as initial and single final state. Hence, by Parikh's Lemma [15],  $CT_k$  is an effectively constructable semilinear set. As  $CT_k^+ = \{c \in CT_k; \forall c \geq 0\}$ , and as systems of linear inequalities are expressible in Presburger Arithmetic the claim of the lemma follows from the fact that semilinear sets are effectively closed under Boolean operations [3]. ■

#### Definition 2:

For  $\bar{m} \in \bar{\mathbf{N}}^n$  and  $N \in \mathbf{N}$  set

$$F(\bar{m}, N) := \{m' \in \mathbf{N}^n; (\forall i \in I_n)[(\bar{m}_i = \omega \wedge m'_i \geq N) \vee (\bar{m}_i = m'_i)]\}.$$

In the sequel, we shall make use of the following basic properties of  $RG(T, m)$  proved in [4,5]:

- a) For any given  $N \in \mathbf{N}$  and node  $k$  in  $RG(T, m)$ , one can effectively find some  $\tau \in T^*$  (and hence  $\Phi(\tau) \in \mathbf{N}^n$ ) s.t.  $a(\tau, m)$  and  $m + \delta\tau \in F(\bar{m}(k), N)$ .
- b) For any given node  $k$  in  $RG(T, m)$ , there is a path from the root  $r$  to  $k$  with edge-labelling sequence  $\tau$  s.t.  $a(\tau, m)$ .

#### Lemma 3:

Let  $(T, m)$  be a persistent VRS and  $k$  a node in  $RG(T, m)$ . Let further  $p$  be the projection of



$\mathbb{N}^n \cup \mathbb{Z}^n$  on those coordinates where  $\bar{m}(k)$  is not equal  $\omega$ . (Note that two transitions  $p(t_i)$  and  $p(t_j)$  with  $i \neq j$  and  $p(t_i) = p(t_j)$  are considered different.) Then  $(p(T), p(\bar{m}(k)))$  is persistent.

*Proof:*

Assume that  $(p(T), p(\bar{m}(k)))$  is not persistent. Then there are  $\tau \in T^*$  and  $t_i, t_j \in T$  with  $i \neq j$  s.t. (with  $m' := \bar{m}(k) + \delta\tau$ )

$$a(p(t_i), p(m')) \wedge a(p(t_j), p(m')) \wedge \neg a(p(t_i)p(t_j), p(m')). \quad (*)$$

Now, effectively find some  $\tau \in T^*$  s.t.  $a(\tau, m)$  and  $m + \delta\tau \in F(\bar{m}(k), N)$ . Choosing  $N$  big enough one could obtain  $m'' \in R(T, m) \cap F(\bar{m}(k), N)$  s.t.  $a(\tau t_i, m'') \wedge a(\tau t_j, m'')$ . From (\*) then follows  $\neg a(\tau t_i t_j, m'')$ , contradicting the persistence of  $(T, m)$ . ■

### Definition 3:

Let  $(T, m)$  be an arbitrary VRS.

- a) A transition  $t \in T$  is *bounded* in  $(T, m)$  iff  $(\exists N \in \mathbb{N}, \nexists c \in C(T, m))[N\Phi(t) \leq c]$ .
- b) A strongly connected component (SCC)  $CC$  in  $RG(T, m)$  is called *distinguished* iff the labels of all edges leaving  $CC$  do not appear as labels of edges within any SCC.

It has been shown in [5] that it is decidable whether  $t \in T$  is bounded. As a matter of fact,  $t \in T$  is not bounded iff it is the label of some edge within some SCC of  $RG(T, m)$ . Let, in the sequel,  $BT(T, m) \subseteq T$  denote the set of bounded transitions in  $(T, m)$ .

If  $(T, m')$  is a persistent VRS, and if  $\tau, \tau' \in T^*$  are sequences s.t.  $a(\tau, m'), a(\tau', m')$ , and  $\delta\tau \geq 0$  then  $\Phi(\tau) \in C(T, m' + \delta\tau)$ . This follows from the fact (proven in [7]) that  $c, c' \in C(T, m')$  implies  $\max\{c, c'\} - c \in C(T, m' + Vc)$  and the observation that  $a(\tau', m' + \delta\tau)$  because of  $\delta\tau \geq 0$ .

Now, let  $(T, m)$  be a persistent VRS,  $k$  a node in some distinguished SCC of  $RG(T, m)$ ,  $k' \neq k$  a node on a cycle through  $k$ , and  $\tau$  the edge-labelling sequence of a cycle through  $k'$  but not  $k$ . Note that the node markings  $\bar{m}(k'')$  have the same set of  $\omega$ -coordinates for all nodes  $k''$  in the SCC of  $k$ . If  $p$  denotes the projection on those coordinates where  $\bar{m}(k)$  is not equal to  $\omega$ , we have  $a(p(\tau), p(\bar{m}(k'')))$  and  $p(\delta\tau) = 0$ . It is also clear from the construction

in Algorithm 1 that  $p(\bar{m}(k)) \in R(p(T), p(\bar{m}(k')))$ . Therefore, using Lemma 3 and the above observation, there must be a path starting from  $k$  with some edge-labelling sequence  $\tau'$  s.t.  $\Phi(\tau') = \Phi(\tau)$ . But then all transitions in  $\tau'$  are unbounded,  $p(\delta\tau') = 0$ , and this path must end in  $k$ . From this observation, one easily obtains

**Lemma 4:**

If  $(T, m)$  is a persistent VRS and  $k$  a node in some distinguished SCC of  $RG(T, m)$  then  $CT_{k'}$  (resp.  $CT_{k'}^+$ ) is linear and equal for all  $k'$  in the SCC of  $k$ .

*Proof:*

From the above discussion, one deduces that the sets  $CT_{k'}$  are linear and equal for all  $k'$  in the SCC of  $k$ . But as  $CT_{k'}^+ = \{c \in CT_{k'}; \forall c \geq 0\}$ , so are the  $CT_{k'}^+$ . ■

On the other hand, one may observe that if  $(T, m)$  is an arbitrary VRS and  $CC$  some distinguished SCC in  $RG(T, m)$  s.t. all  $\bar{m}(k)$  for  $k$  in  $CC$  are persistent, then  $CT_k$  (resp.  $CT_k^+$ ) is linear and equal for all  $k$  in  $CC$  because  $\{\bar{m}(k); k \text{ in } CC\} = R(T - BT(T, m), \bar{m}(k'))$  for any  $k'$  in  $CC$ , and  $(T - BT(T, m), \bar{m}(k'))$  still is persistent.

The following theorem states the basic properties of distinguished SCC's in the reachability graph of persistent VRS's. Let, in the sequel,  $w' := |BT(T, m)|$  denote the cardinality of  $BT(T, m) \subseteq T$ , and  $p_{BT}$  the projection of  $\mathbb{N}^w$  onto those coordinates which correspond to transitions in  $BT$ .

**Theorem 1:**

Let  $(T, m)$  be a persistent VRS.

- a) There is exactly one maximal SCC (i.e. no other SCC can be reached from it) in  $RG(T, m)$ .
- b) For each  $\bar{c} \in \{p_{BT}(c); c \in CT(T, m)\}$ , there is exactly one distinguished SCC  $CC(\bar{c})$  in  $RG(T, m)$  s.t.  $(\forall \tau \in T^*)[(a(\tau, m) \wedge \tau \text{ determines the edge-labelling sequence of some path$

in  $RG(T, m)$  from the root  $r$  to some node in  $CC(\bar{c}) \Rightarrow p_{BT}(\Phi(\tau)) = \bar{c}$ .

*Proof:*

a) Assume that there are two nodes  $k$  and  $k'$  in two different maximal SCC's of  $RG(T, m)$ . Then there are  $\tau, \tau' \in T^*$  s.t.  $a(\tau, m), a(\tau', m)$ , and  $\tau$  (resp.,  $\tau'$ ) determine a path in  $RG(T, m)$  from the root to  $k$  (resp.,  $k'$ ). Because of the result in [7] and its consequence mentioned above, we may assume w.l.g. that  $\Phi(\tau) = \Phi(\tau')$  and  $\bar{m}(k)$  and  $\bar{m}(k')$  have the same set of  $\omega$ -coordinates, and hence, that  $k = k'$ , contradicting the assumption.

b) First note that in  $RG(T, m)$  an edge the label of which is a bounded transition always leads from one SCC to a different one.

Let  $\bar{c} \in \{p_{BT}(c); c \in C(T, m)\} \subseteq \mathbf{N}^{w'}$ . Obtain from  $(T, m)$  a modified  $(T', m')$  in the following way (where  $BT = \{t_{i_1}, \dots, t_{i_{w'}}\}$ ):

Add to all  $(u_i, v_i) \in T$   $w'$  new coordinates the  $j$ -th of which is 1 for  $u_i$ ,  $-1$  for  $v_i$ , for  $j \in I_{w'}$ , and zero in all other cases, and  $m'$  equals  $(m, \bar{c}) \in \mathbf{N}^{w+w'}$ . It is easy to see from the definition that  $(T', m')$  still is persistent, and that  $C(T', m') = \{c \in C(T, m); p_{BT}(c) \leq \bar{c}\}$ . Also, it follows from the remark made above that for all  $\tau' \in T'^*$  with  $a(\tau', m') p_{BT}(\Phi(\tau')) = \bar{c}$  if  $\tau'$  determines a path in  $RG(T', m')$  from the root to some node in the maximal SCC.

If one now observes that—again because of the above remark— $RG(T', m')$  is isomorphic to a subgraph of (some, because Algorithm 1 is nondeterministic)  $RG(T, m)$  (with the canonical mapping between the edge and node labels, resp.), then let the maximal SCC in  $RG(T', m')$  correspond to  $CC(\bar{c})$  in  $RG(T, m)$ , and b) follows from a). ■

#### Definition 4:

Let  $(T, m)$  be a VRS and  $k$  a node in  $RG(T, m)$  s.t.  $CT_k^+$  is linear. Let, further,  $\tau^j, j \in I_h$ , be edge-labelling sequences of paths in  $RG(T, m)$  from  $k$  to  $k$  s.t.

$$CT_k^+ = \left\{ \sum_{j=1}^h n_j \Phi(\tau^j); (n_1, \dots, n_h) \in \mathbf{N}^h \right\}.$$

A *hurdle* for  $k$  is then a number  $H_k \in \mathbf{N}$  s.t.

$$m' \in F(\bar{m}(k), H_k) \Rightarrow (\forall j \in I_h)[a(\tau^j, m')].$$

Given  $RG(T, m)$  for some  $(T, m)$ , an  $H_k$  for some  $k$  with linear  $CT_k^+$  can effectively be

determined.

Finally, we note a property of linear sets which is used by the algorithms discussed in the next section. Let, for sets  $A, B \subseteq \mathbb{N}^n$ ,  $A + B$  denote the set  $\{a + b; a \in A, b \in B\}$ . Then we have

**Lemma 5:**

Let  $L \subseteq \mathbb{N}^n$  be linear,  $0 \in L$ , and  $L'$  some subset of  $L$ . Then there is a finite  $B \subseteq L'$  s.t.  $L' \subseteq B + L$ . If  $L'$  is semilinear, and  $L, L'$  are effectively given, such a  $B$  can effectively be obtained.

The proof is left to the reader.

#### 4. A decision procedure for persistence

Let  $k$  be a node in the reachability graph  $RG(T, m)$  of some VRS  $(T, m)$ ,  $m \in \mathbb{N}^v$ , s.t.  $CT_{k'}$  is equal and linear for all nodes  $k'$  in the SCC of  $k$ . We are now going to describe a procedure *slset* which for SCC's as above constructs a semilinear representation of the set  $\{c + \Phi(\tau); a(\tau, m + Vc) \text{ and } \tau \text{ is the edge-labelling sequence of a path in the SCC of } k\}$  where  $c$  is some counter s.t.  $m + Vc \in F(\bar{m}(k), H_k)$ .

```

procedure slset (k:node; SL:repr of semilinear set; c:counter):
begin
  co it is required that  $k$  is a node in the reachability graph  $RG(T, m)$  of some VRS  $(T, m)$ ,  $m \in \mathbb{N}^v$ , s.t.  $CT_{k'}$  is
    equal and linear for all  $k'$  in the SCC of  $k$ .  $(T, m)$  etc. are global for slset oc;
  var HK:integer; CTK:repr of semilinear set;
  co SL, CTK refer to representations of semilinear sets oc;
  procedure complete (k:node; c:counter);
  begin
    var i:integer; bset:finite set of counter; L':repr of semilinear set;
    for  $i := 1, \dots, w$  do
      if there is an edge labelled  $t_i$  from  $k$  to some  $k'$  (possibly  $k = k'$ ) in the SCC of  $k$  s.t.
         $(\exists c' \in c + CTK)[a(t_i, m + Vc') \wedge (c' + \Phi(t_i) \notin SL)]$ 
        co this can be written as a Presburger expression oc
      then
         $L' := \{c' \in c + CTK; a(t_i, m + Vc') \wedge (c' + \Phi(t_i) \notin SL)\}$ ;
        co the right hand side defines a semilinear set oc;
         $bset :=$  some finite subset of  $L'$  s.t.  $L' \subseteq bset + CTK$ ;
        co because of Lemma 5, this assignment is effective oc;
        for all  $c'$  in bset do
           $SL := SL \cup (c' + \Phi(t_i) + CTK)$ ;
          complete( $k'$ ,  $c' + \Phi(t_i)$ )
        od
      fi
    od
  end complete;
   $CTK :=$  co a representation of oc  $CT_k^+$ ;
  co note Lemma 2 and the remark after Lemma 4 oc;
   $HK :=$  some hurdle  $H_k$  for  $k$ ;
   $c :=$  some counter  $\in C(T, m)$  s.t.  $m + Vc \in F(\bar{m}(k), HK)$ ;
   $SL :=$  co a representation of oc  $c + CT_k^+$ ;
  complete( $k$ ,  $c$ )
end slset;

```

**Theorem 2:**

- a) The procedure *slset* terminates and yields a set  $SL \subseteq c(T, m)$ .
- b) If  $(T, m)$  is persistent then  $SL = \{c + \Phi(\tau); \tau \text{ is the edge-labelling sequence of a path in the SCC of } k \text{ in } RG(T, m) \text{ starting from } k \text{ s.t. } a(\tau, m + Vc)\}$ , where  $c$  is the counter determined in *slset*.

*Proof:*

a) From the selection of  $HK$  and  $c$ , it is clear that  $c + CT_k^+ \subseteq C(T, m)$ . When some  $c' + \Phi(t_i) + CTK$  is added to  $SL$  one can assume by induction on the depth of recursion that  $SL \subseteq c + C(T, m + Vc)$  before that step. Let  $\tau' \in T^*$  be some sequence s.t.  $a(\tau', m + Vc) \wedge \Phi(\tau') = c' - c$ . Furthermore,  $SL + CTK \subseteq SL$ . But by the definition of  $HK$  and  $CTK$ , there is, for any  $c'' \in CTK$ , a  $\tau'' \in T^*$  s.t.  $a(\tau'', m + Vc) \wedge \Phi(\tau'') = c'' \wedge \tau''$  determines a path from  $k$  to  $k$ . Hence,  $a(\tau', m + V(c + c'')) \wedge a(t_i, m + V(c' + c''))$  as  $\delta\tau'' = Vc'' \geq 0$ . As  $c''$  has been chosen arbitrary in  $CTK$  this shows that  $c' + \Phi(t_i) + CTK \subseteq c + C(T, m + Vc)$ , and by induction, that invariantly  $SL \subseteq c + C(T, m + Vc) \subseteq C(T, m)$ . Now assume that *slset* does not terminate. Then, by König's Infinity Lemma, there must be an infinite chain of nested recursive calls of the procedure *complete* and a subchain of this chain such that all calls in this subchain have the same first parameter. Let  $(c^i)_{i \in \mathbb{N}}$  be the sequence of counters in the second parameter position of this subchain. Because each infinite sequence in  $\mathbb{N}^e$  has a nondecreasing infinite subsequence (this is a corollary of what is sometimes referred to as Dickson's Lemma [2, Lemma A]) there is a subsequence  $(c^i)_{i \in \mathbb{N}}$  of  $(c^i)_{i \in \mathbb{N}}$  s.t.  $(m + Vc^i)_{i \in \mathbb{N}}$  is nondecreasing. As has been shown above, for each  $c^i$  there is a  $\tau^i \in T^*$  s.t.  $a(\tau^i, m + Vc) \wedge \Phi(\tau^i) = c^i - c \wedge (\tau^i \text{ is the edge-labelling sequence of a path } \alpha^i \text{ in the SCC of } k \text{ in } RG(T, m) \text{ starting from } k \text{ and ending in some fixed node } k')$ . The last observation follows from the choice of  $(c^i)_{i \in \mathbb{N}}$ . Considering the multiplicity with which the edges of  $RG(T, m)$  appear in  $\alpha^i$ ,  $i \in \mathbb{N}$ , and applying once more Dickson's Lemma, one obtains indices  $j < j'$  s.t.  $\alpha^{j'}$  contains each edge of  $RG(T, m)$  at least as often as does  $\alpha^j$ . As  $CT_{k'}^+$  is linear and equal for all  $k'$  in the SCC of  $k$ , and by the definition of the  $\tau^i$ ,  $\Phi(\tau^{j'}) - \Phi(\tau^j) \in CTK$ . Thus, after  $\bar{c}^j + CTK$  has been added to  $SL$ ,  $\bar{c}^j \in SL$ , contradicting the assumption that *slset* does not terminate.

b) From the first part of this proof we know that  $SL \subseteq \{c + \Phi(\tau); \tau \text{ is the edge-labelling sequence of a path in the SCC of } k \text{ in } RG(T, m) \text{ starting from } k \text{ s.t. } a(\tau, m + Vc)\}$ . The other direction can easily be seen by induction on the length of  $\tau$ . ■

The procedure *sset* will now be used in the following main algorithm to decide persistence of an arbitrary VRS.

### Algorithm 2:

**begin**

var *GSL, NPC*:repr of semilinear set;

procedure test ((*T, m*):VRS);

**begin**

var *i*:integer; *SL, SL'*:repr of semilinear set; *k*:node; *c', cmax*:counter;

procedure *sset* (*k*:node; ... ); ... ;

construct *RG(T, m)* using Algorithm 1;

if *RG(T, m)* doesn't satisfy the necessary condition of Theorem 1a) or contains a non-persistent node marking thus violating Lemma 3

then stop '(*T, m*) is not persistent' **fi**;

set *CC<sub>i</sub>*, *i* = 1, ..., *h*, the distinguished SCC's of *RG(T, m)*;

*cmax* := 0 co  $\in \mathbb{N}^w$  **oc**;

*SL* :=  $\emptyset$ ;

for *i* := 1, ..., *h* do

*k* := some node in *CC<sub>i</sub>*;

*sset*(*k, SL', c'*);

*cmax* := max(*c', cmax*) co max component-wise **oc**;

*SL* := *SL*  $\cup$  *SL'*

**od**;

if *SL*  $\cap$  *NPC*  $\neq \emptyset$  then stop '(*T, m*) is not persistent' **fi**;

*GSL* := *GSL*  $\cup$  *SL*;

co *GSL* globally collects all counters in *C(T, m)* found by the algorithm **oc**;

for *i* := 1, ..., *w* do

    if *t<sub>i</sub>*  $\in T - BT(T, m)$  then

        construct from (*T, m*) a new VRS (*T<sup>i</sup>, m<sup>i</sup>*) where *T<sup>i</sup>* is obtained from *T* by adding a *w* + 1st coordinate which is 1 for *u<sub>i</sub>*, -1 for *v<sub>i</sub>*, and zero in all other cases, and *m<sup>i</sup>* equals (*m, cmax<sub>i</sub>*)  $\in \mathbb{N}^{w+1}$ ;

        co this means that  $C(T^i, m^i) = \{c \in C(T, m); c_i \leq cmax_i\}$  **oc**;

        test((*T<sup>i</sup>, m<sup>i</sup>*));

        co note that *GSL* is global in this recursion **oc**

**fi**

**od**

**end test**;

*NPC* := *NPC(T, m)* co note Lemma 1 **oc**;

*GSL* :=  $\emptyset$ ;

test((*T, m*));

if ( $\exists c \in GSL, \exists i \in I_w$ ) [*a(t<sub>i</sub>, m + Vc)*  $\wedge$  (*c +  $\Phi(t<sub>i</sub>)$*   $\notin GSL$ )]

then stop '(*T, m*) is not persistent'

else stop '(*T, m*) is persistent'

**fi**

**end Algorithm 2.**

**Theorem 3:**

Algorithm 2 terminates for every VRS  $(T, m)$  and determines whether  $(T, m)$  is persistent or not.

**Corollary 1:**

Persistence is decidable for arbitrary VRS's.

**Corollary 2:**

If  $(T, m)$  is persistent Algorithm 2 yields  $GSL$  s.t.  $GSL = C(T, m)$ .

**Corollary 3:**

There is an effective construction of semilinear representations of the reachability set of persistent VRS's.

*Proof:*

Because of the reduction of the number of unbounded transitions in successive recursive calls of the procedure *test*, it is clear that Algorithm 2 terminates. If Algorithm 2 stops within *test* the answer given is correct because of Lemma 3. Otherwise, if the condition

$$(\exists c \in GSL, \exists i \in I_w)[a(t_i, m + Vc) \wedge (c + \Phi(t_i) \notin GSL)]$$

in the last **if**-statement of Algorithm 2 evaluates to **false**,  $GSL$  equals  $C(T, m)$  because of Theorem 2a), and  $(T, m)$  is persistent because, in fact,  $GSL \cap NPC = \emptyset$  has been verified by the algorithm. Conversely, if  $(T, m)$  is persistent then so are all  $(T^i, m^i)$  generated in the recursive calls of *test* as addition of a coordinate just bounding the number of times how often some transition can be applied doesn't hurt persistence. Furthermore, if  $c \in C(T, m)$ , then it follows from Theorem 1b), Theorem 2b), and Lemma 3.1 in [9] that

$$((\forall i \in I_w)[t_i \in T - BT(T, m) \Rightarrow c_i \geq \max_i] \Rightarrow c \in GSL).$$

Hence, by induction on the number of unbounded transitions and by the construction of the  $(T^i, m^i)$ ,  $C(T, m) \subseteq GSL$ , and because of Theorem 2a),  $C(T, m) = GSL$ . Therefore, if  $(T, m)$  is persistent, the condition in the last **if**-statement of Algorithm 2 evaluates to **false**. This proves the theorem. The corollaries are immediate consequences. ■



## 5. Conclusion

The algorithm presented in this report extends the results in [11,12] and answers some of the questions asked in [9]. Thus, persistence of arbitrary VRS's is decidable (by an algorithm which does not make use of a solution to the general reachability problem), and there is an effective method to construct semilinear representations for the reachability set of persistent VRS's. As far as the author knows,  $m$ -reversible [1] and persistent VRS's are the only classes of VRS's for which an effective representation of infinite reachability sets has been given so far. Thus there remains a number of open problems in extending this result to other classes of VRS's of interest as well as in establishing the complexity of the algorithm given here. Another open problem concerns the characterization of the class of VRS's that have semilinear reachability sets.

**Addendum:** While this report was being prepared, another algorithm for deciding persistence based on the non-constructive proof in [9] was obtained independently in [18].

## 6. References

1. Araki, T., Kasami, T.:  
Decidable Problems on the Strong Connectivity of Petri Net Reachability Sets.  
Theoret. Comp. Sci. 4, 1 (Febr. 1977), p.99-119
2. Dickson, L.E.:  
Finiteness of the Odd Perfect and Primitive Abundant Numbers with  $n$  Distinct Prime Factors.  
Amer. J. Math. 35 (1913), p.413-422
3. Ginsburg, S., Spanier, E.H.:  
Semigroups, Presburger Formulas, and Languages.  
Pacific J. of Math. 16 (1966), p.285-296
4. Hack, M.:  
Decision Problems for Petri Nets and Vector Addition Systems.  
M.I.T., Project MAC, MAC-TM 59 (March 1975)
5. Hack, M.:  
Decidability Questions for Petri Nets.  
M.I.T., LCS, TR 161 (June 1976)
6. Karp, R.M., Miller, R.E.:  
Parallel Program Schemata.  
JCSS 3, 2 (May 1969), p.147-195
7. Keller, R.M.:  
A Fundamental Theorem of Asynchronous Parallel Computation.  
In: Feng, T.Y.(ed.): Parallel Processing, Proc. Sagamore Computer Conference. LNCS 24. Springer 1975, p.102-112
8. Keller, R.M.:  
Vector Replacement Systems: A Formalism for Modelling Asynchronous Systems.  
Princeton University, CSL, TR 117 (Dec. 1972)
9. Landweber, L.H., Robertson, E.L.:  
Properties of Conflict Free and Persistent Petri Nets.  
JACM 25 (1978), p.352-364
10. Lipton, R.J., et al.:  
Synchronization and Computing Capabilities of Linear Asynchronous Structures.  
Proc. 16th Ann. Symp. on FOCS (1975), p.19-28
11. Mayr, E.W.:  
An Effective Representation of the Reachability Set of Persistent Petri Nets.  
M.I.T., LCS, TM 188
12. Müller, H.:  
Decidability of Reachability in Persistent Vector Replacement Systems.  
Proc. 9th Symp. on MFCS (1980). LNCS 88. Springer 1980, p.426-438
13. Muller, D.E., Bartky, M.S.:  
A Theory of Asynchronous Circuits.  
Proc. Int. Symp. on Theory of Switching, Harvard University Press (Cambridge, Mass.) (1959), p.204-243
14. Oppen, D.C.:  
A  $2^{2^{2^n}}$  Upper Bound on the Complexity of Presburger Arithmetic.  
JCSS 16 (1978), p.323-332
15. Parikh, R.J.:  
On Context-free Languages.  
JACM 13 (1966), p.570-581

16. Presburger, M.:  
Über die Vollständigkeit eines gewissen Systems der Arithmetik ganzer Zahlen, in welchem die Addition als einzige Operation hervortritt.  
Compte-Rendus du I. Congrès des Mathématiciens des pays Slavs, Warsaw (1930), p.92-101
17. Rosen, B.:  
Tree Manipulating Systems and Church-Rosser Theorems.  
JACM 20 (1973), p.160-187
18. Grabowski, J.:  
The Decidability of Persistence for Vector Addition Systems.  
IPL 11, 1 (1980), p.20-23